

## Farey Fractions

### Definition

The set of Farey fractions of order  $n$ , denoted by  $F_n$ , is the set of reduced fractions in the closed interval  $[0, 1]$  with denominators not exceeding  $n$  listed in increasing order.

### Examples

$F_1$	$\frac{0}{1}$																		$\frac{1}{1}$
$F_2$	$\frac{0}{1}$								$\frac{1}{2}$										$\frac{1}{1}$
$F_3$	$\frac{0}{1}$						$\frac{1}{3}$		$\frac{1}{2}$		$\frac{2}{3}$								$\frac{1}{1}$
$F_4$	$\frac{0}{1}$			$\frac{1}{4}$		$\frac{1}{3}$		$\frac{1}{2}$		$\frac{2}{3}$		$\frac{3}{4}$							$\frac{1}{1}$
$F_5$	$\frac{0}{1}$		$\frac{1}{5}$	$\frac{1}{4}$		$\frac{1}{3}$	$\frac{2}{5}$		$\frac{1}{2}$		$\frac{3}{5}$	$\frac{2}{3}$		$\frac{3}{4}$	$\frac{4}{5}$				$\frac{1}{1}$
$F_6$	$\frac{0}{1}$	$\frac{1}{6}$	$\frac{1}{5}$	$\frac{1}{4}$		$\frac{1}{3}$	$\frac{2}{5}$		$\frac{1}{2}$		$\frac{3}{5}$	$\frac{2}{3}$		$\frac{3}{4}$	$\frac{4}{5}$	$\frac{5}{6}$			$\frac{1}{1}$
$F_7$	$\frac{0}{1}$	$\frac{1}{7}$	$\frac{1}{6}$	$\frac{1}{5}$	$\frac{1}{4}$	$\frac{2}{7}$	$\frac{1}{3}$	$\frac{2}{5}$	$\frac{3}{7}$	$\frac{1}{2}$	$\frac{4}{7}$	$\frac{3}{5}$	$\frac{2}{3}$	$\frac{5}{7}$	$\frac{3}{4}$	$\frac{4}{5}$	$\frac{5}{6}$	$\frac{6}{7}$	$\frac{1}{1}$

### Remark

These examples illustrate some general properties of Farey fractions. For example,  $F_n \subset F_{n+1}$ , so we get  $F_{n+1}$  by inserting new fractions in  $F_n$ . If  $\frac{a}{b} < \frac{c}{d}$  are consecutive in  $F_n$  and separated in  $F_{n+1}$ , then the fraction  $\frac{a+c}{b+d}$  does the separating, and no new ones are inserted between  $\frac{a}{b}$  and  $\frac{c}{d}$ .

This new fraction is called the *mediant* of  $\frac{a}{b}$  and  $\frac{c}{d}$ .

### Theorem 1

If  $\frac{a}{b} < \frac{c}{d}$ , their mediant  $\frac{a+c}{b+d}$  lies between them.

PROOF

$$\frac{a+c}{b+d} - \frac{a}{b} = \frac{bc - ad}{b(b+d)} > 0 \quad \text{and} \quad \frac{c}{d} - \frac{a+c}{b+d} = \frac{bc - ad}{d(b+d)} > 0 .$$

**Theorem 2**

Let  $0 \leq \frac{a}{b} < \frac{c}{d} \leq 1$  with  $bc - ad = 1$ .

Then  $\frac{a}{b}$  and  $\frac{c}{d}$  are consecutive terms in  $F_n$  for the following values of  $n$  :

$$\max \{b, d\} \leq n \leq b + d - 1 .$$

PROOF

The condition  $bc - ad = 1$  implies that the fractions  $\frac{a}{b}$  and  $\frac{c}{d}$  are reduced.

If  $\max \{b, d\} \leq n$  then  $b \leq n$  and  $d \leq n$  so  $\frac{a}{b}$  and  $\frac{c}{d}$  are certainly in  $F_n$ .

Now we prove that they are consecutive if  $n \leq b + d - 1$ .

If they are not consecutive, then there is another fraction  $\frac{h}{k}$  between them, that is  $\frac{a}{b} < \frac{h}{k} < \frac{c}{d}$ .

Since

$$k = k(bc - ad) = b(ck - dh) + d(bh - ak) ,$$

and

$$\frac{a}{b} < \frac{h}{k} < \frac{c}{d} \text{ implies that } ck - dh \geq 1 \text{ and } bh - ak \geq 1 ,$$

then

$$k = b(ck - dh) + d(bh - ak) \geq b + d .$$

Therefore, if  $n \leq b + d - 1$ , then  $\frac{a}{b}$  and  $\frac{c}{d}$  must be consecutive in  $F_n$ .

**Theorem 3**

Let  $0 \leq \frac{a}{b} < \frac{c}{d} \leq 1$  with  $bc - ad = 1$ .

If  $\frac{h}{k}$  is the mediant of  $\frac{a}{b}$  and  $\frac{c}{d}$ , then  $\frac{a}{b} < \frac{h}{k} < \frac{c}{d}$  and  $bh - ak = 1$ ,  $ck - dh = 1$ .

PROOF

Since  $\frac{h}{k}$  lies between  $\frac{a}{b}$  and  $\frac{c}{d}$ , then  $ck - dh \geq 1$  and  $bh - ak \geq 1$ . Thus

$$k = b(ck - dh) + d(bh - ak)$$

shows that  $k = b + d$  if and only if  $ck - dh = 1$ ,  $bh - ak = 1$ .

**Theorem 4**

The set  $F_{n+1}$  includes  $F_n$ .

Each fraction in  $F_{n+1}$  which is not in  $F_n$  is the mediant of a pair of consecutive fractions in  $F_n$ .

Moreover, if  $\frac{a}{b} < \frac{c}{d}$  are consecutive in any  $F_n$ , then they satisfy the unimodular relation  $bc - ad = 1$ .

PROOF

Use induction on  $n$ .

When  $n = 1$ , the fractions  $\frac{0}{1}$  and  $\frac{1}{1}$  are consecutive and satisfy the unimodular relation.

We pass from  $F_1$  to  $F_2$  by inserting  $\frac{1}{2}$ .

Now suppose  $\frac{a}{b}$  and  $\frac{c}{d}$  are consecutive in  $F_n$  and satisfy the unimodular relation  $bc - ad = 1$ .

Then by Theorem 2, they will be consecutive in  $F_m$  for all  $m$  satisfying

$$\max \{b, d\} \leq m \leq b + d - 1 .$$

Form their mediant  $\frac{h}{k}$  where  $h = a + c$ ,  $k = b + d$ .

By Theorem 3,  $bh - ak = 1$ ,  $ck - dh = 1$ , so  $h$  and  $k$  are relatively prime.

The fractions  $\frac{a}{b}$  and  $\frac{c}{d}$  are consecutive in  $F_m$  for all  $m$  satisfying  $\max \{b, d\} \leq m \leq b + d - 1$ , but are not consecutive in  $F_k$  since  $k = b + d$  and  $\frac{h}{k}$  lies in  $F_k$  between  $\frac{a}{b}$  and  $\frac{c}{d}$ .

But the two new pairs  $\frac{a}{b} < \frac{h}{k}$  and  $\frac{h}{k} < \frac{c}{d}$  are now consecutive in  $F_k$  because

$$k = \max \{b, k\} \text{ and } k = \max \{d, k\}.$$

The two new pairs still satisfy the unimodular relations  $bh - ak = 1$ ,  $ck - dh = 1$ .

This shows that in passing from  $F_n$  to  $F_{n+1}$  every new fraction inserted must be the mediant of a consecutive pair in  $F_n$  and the new consecutive pairs satisfy the unimodular relations. Therefore  $F_{n+1}$  has these properties if  $F_n$  does.

**Theorem 5**

If  $\frac{a}{b}$  and  $\frac{c}{d}$  are consecutive in  $F_n$ , then

$$\left| \frac{a}{b} - \frac{a+c}{b+d} \right| = \frac{1}{b(b+d)} \leq \frac{1}{b(n+1)}$$

and

$$\left| \frac{c}{d} - \frac{a+c}{b+d} \right| = \frac{1}{d(b+d)} \leq \frac{1}{d(n+1)}.$$

**Theorem 6**

Given any real number  $\theta$  and any positive integer  $N$ , there exist rational numbers  $\frac{a}{b}$  such that

$$\left| \theta - \frac{a}{b} \right| < \frac{1}{b(N+1)} \quad \text{with } 0 < b \leq N.$$

**Theorem 7**

Given any irrational  $\theta$ , there exist infinitely many rational numbers  $\frac{a}{b}$  such that

$$\left| \theta - \frac{a}{b} \right| < \frac{1}{b^2}.$$

**Theorem 8**

If  $m$  and  $n$  are positive integers then not both of the inequalities can hold:

$$\frac{1}{mn} \geq \frac{1}{\sqrt{5}} \left( \frac{1}{m^2} + \frac{1}{n^2} \right)$$

and

$$\frac{1}{m(m+n)} \geq \frac{1}{\sqrt{5}} \left( \frac{1}{m^2} + \frac{1}{(m+n)^2} \right).$$

**Theorem 9     Hurwitz**

Given any irrational  $\theta$ , there are infinitely many rational numbers  $\frac{a}{b}$  such that

$$\left| \theta - \frac{a}{b} \right| < \frac{1}{\sqrt{5} a^2}.$$

Moreover the result is false if  $\frac{1}{\sqrt{5}}$  is replaced by any smaller constant.